

Note di Matematica

Note Mat. 30 (2010) n. 1, 107–109.

ISSN 1123-2536, e-ISSN 1590-0932

doi:10.1285/i15900932v30n1p107

# On angular bisectors in normed linear spaces

**Horst Martini**

Faculty of Mathematics, Chemnitz University of Technology

horst.martini@mathematik.tu-chemnitz.de

**Senlin Wu**<sup>i</sup>

Department of Applied Mathematics, Harbin University of Science and Technology

senlin.wu@hotmail.com

Received: 29.04.2009; accepted: 20.10.2009.

**Abstract.** We prove that a Minkowski space is Euclidean if it has the weak bisector property. This confirms a conjecture of R. W. Freese, C. R. Diminnie, and E. Z. Andalafte.

**Keywords:** angular bisector, angular measure, characterization of inner product spaces, Minkowski plane, Minkowski space, norm, weak bisector property

**MSC 2000 classification:** primary 46B20, secondary 46C15, 52A21

By  $X$  we denote the *Minkowski space* (i.e., the finite dimensional real Banach space) with origin  $o$ , norm  $\|\cdot\|$ , and *unit sphere*  $S_X := \{x \in X : \|x\| = 1\}$ . When  $X$  is a Minkowski plane, i.e., a two-dimensional Minkowski space,  $S_X$  is also called the *unit circle* of  $X$ . Basic references for the geometry of Minkowski spaces are [5], [4], and the monograph [8]. For  $x \neq -y$ , the intersection of the cone  $\{\lambda x + \mu y : \lambda, \mu \geq 0\}$  and  $S_X$  is called the (*undirected*) *arc* between  $x$  and  $y$  and denoted by  $S_X(x, y)$ , and the length of  $S_X(x, y)$  is denoted by  $\delta_X(x, y)$ . For brevity, we use the shorthand notation  $\hat{x} = \frac{x}{\|x\|}$  for any point  $x \neq o$ .

For any three non-collinear points  $p, x, y \in X$  we call the convex set bounded by the rays  $[p, x)$  and  $[p, y)$  the *angle*  $xpy$  (denoted by  $\angle xpy$ ) with  $p$  as apex. For two linearly independent points  $x, y \in S_X$  the authors of [2] defined the measure of  $\angle xoy$  by

$$A(x, y) := \cos^{-1} \left[ \frac{1}{2} (2 - \|\hat{x} - \hat{y}\|^2) \right].$$

The ray  $[o, z)$  with  $z \in \angle xoy$  is called the *angular bisector* of  $\angle xoy$  provided  $A(x, z) = A(y, z)$  (the uniqueness of  $z$  in this framework follows from [3, p. 170, Corollary]). R. W. Freese, C. R. Diminnie, and E. Z. Andalafte in [3] proved that a Minkowski space  $X$  is Euclidean if  $X$  has the *angle bisector property*, where  $X$  is said to have the angle bisector property if for all linearly independent  $x, y \in S_X$  the point  $z = \widehat{x+y}$  satisfies  $A(x, z) = A(y, z) = \frac{1}{2} A(x, y)$ . In [6] a stronger result was proved (see Theorem 4.2 there): If for all linearly independent  $x, y \in S_X$  the point  $z = \widehat{x+y}$  satisfies  $A(x, z) = A(y, z)$ , then the underlying Minkowski plane is Euclidean. The authors of [3] also asked whether the *weak bisector property*, which is formulated in the following, still implies that the underlying space is Euclidean.

<sup>i</sup>The research of the second named author is supported by a grant from the Harbin University of Science and Technology (grant number 2009YF028) and by Deutsche Forschungsgemeinschaft.




Figure 1. Proof of Theorem 1.

**Weak bisector property:** For any two linearly independent points  $x, y \in S_X$  there exists a point  $z \in \angle xoy$  having the property  $A(x, z) = A(y, z)$  which satisfies  $A(x, z) = \frac{1}{2}A(x, y)$ .

Once more we mention that in view of [3, p. 170, Corollary] this point  $z$  is unique, up to positive multiples. The aim of this paper is to give an affirmative answer to this question, namely by

**Theorem 1.** A Minkowski space  $X$  of dimension at least 2 is Euclidean if it has the weak bisector property.

To prove this result we need the following lemma:

**Lemma 1** (See Corollary 2.5 in [7]). Let  $X$  be a Minkowski plane. If there exists a function  $\varphi : [0, 2] \rightarrow [0, 4]$  such that for any  $u, v \in S_X$  we have  $\delta_X(u, v) = \varphi(\|u - v\|)$ , then  $X$  is Euclidean.

**Proof of Theorem 1:** Since a Minkowski space of dimension at least 2 is Euclidean if and only if each of its two-dimensional subspaces is Euclidean (see [1]), we may assume, without loss of generality, that  $X$  is a Minkowski plane. By Lemma 2, we only need to show that the length of any undirected arc of  $S_X$  is determined by the length of the corresponding chord.

Note first that for any  $x, y \in S_X$  there exists a unique point  $z \in S_X$  with  $z \in \angle xoy$  and  $A(x, z) = A(y, z)$ , and that  $\|x - z\| = \|y - z\|$  is determined only by  $\|x - y\|$ . Indeed, from the assumption of the theorem it follows that

$$\cos^{-1} \left[ \frac{1}{2}(2 - \|\hat{x} - \hat{z}\|^2) \right] = A(x, z) = \frac{1}{2}A(x, y) = \frac{1}{2} \cos^{-1} \left[ \frac{1}{2}(2 - \|\hat{x} - \hat{y}\|^2) \right],$$

which implies that

$$\|\hat{x} - \hat{z}\| = \sqrt{2 - \sqrt{4 - \|\hat{x} - \hat{y}\|^2}}.$$

Now, for any points  $x, y, x', y' \in S_X$  with  $\|x - y\| = \|x' - y'\|$ , we show that  $\delta_X(x, y) = \delta_X(x', y')$ . For any integer  $n \geq 1$  we have two subsets  $\{z_1, \dots, z_{2n+1}\}, \{z'_1, \dots, z'_{2n+1}\} \subset S_X$

such that the following holds (note that the points  $z_i$  etc. are found again in view of [3, p. 170, Corollary]):

- (1)  $\{z_1, \dots, z_{2^n+1}\} \subset S_X(x, y)$  and  $\{z'_1, \dots, z'_{2^n+1}\} \subset S_X(x', y')$ ,
- (2)  $A(z_1, z_{2^n+1}) = A(z_{2^n+1}, z_{2^n+1}) = A(z'_1, z'_{2^n+1}) = A(z'_{2^n+1}, z'_{2^n+1})$ ,
- (3)

$$\begin{aligned} A(z_1, z_{2^n+1}) &= A(z_{2^n+1}, z_{2^n+1}) \\ &= A(z'_1, z'_{2^n+1}) = A(z'_{2^n+1}, z'_{2^n+1}) \end{aligned}$$

and

$$\begin{aligned} A(z_{2^n+1}, z_{2^n+1}) &= A(z_{2^n+1}, z_{2^n+1}) \\ &= A(z'_{2^n+1}, z'_{2^n+1}) \\ &= A(z'_{2^n+1}, z'_{2^n+1}) \end{aligned}$$

...

Continuing with similar arguments, we finally obtain that

$$\|z_i - z_{i+1}\| = \|z_j - z_{j+1}\| = \|z'_i - z'_{i+1}\| = \|z'_j - z'_{j+1}\|$$

holds for any  $1 \leq i, j \leq 2^n$ .

Hence

$$\delta_X(x, y) = \sup_{n \geq 1} \left\{ \sum_{i=1}^{2^n} \|z_i - z_{i+1}\| \right\} = \sup_{n \geq 1} \left\{ \sum_{i=1}^{2^n} \|z'_i - z'_{i+1}\| \right\} = \delta_X(x', y'),$$

and this supremum of sums converges to the arc length since  $\|z_i - z_{i+1}\|$  converge to zero. This follows since that sum is bounded from above by the arc length between  $x$  and  $y$ , and we have  $2^n$  summands for  $n \rightarrow \infty$ . The proof is complete.  $\square$

**Remark 1.** The weak bisector property in Theorem 1 can be replaced by the following condition: There exists a real function  $\phi$  such that for any two linearly independent points  $x, y \in S_X$ , the point  $z \in \angle xoy$  having the property  $A(x, z) = A(y, z)$  satisfies  $A(x, z) = \phi(A(x, y))$ .

## References

- [1] D. AMIR, Characterizations of Inner Product Spaces, Birkhäuser, Basel, 1986.
- [2] C. R. DIMINNIE, E. Z. ANDALAFTE, R. W. FREESE, *Angles in normed linear spaces and a characterization of real inner product spaces*, Math. Nachr., **129** (1986), 197–204.
- [3] R. W. FREESE, C. R. DIMINNIE, E. Z. ANDALAFTE, *Angle bisectors in normed linear spaces*, Math. Nachr., **131** (1987), 167–173.
- [4] H. MARTINI, K. J. SWANEPOEL, *The geometry of Minkowski spaces - a survey. Part II*, Expositiones Math., **22** (2004), 93–144.
- [5] H. MARTINI, K. J. SWANEPOEL, G. WEISS, *The geometry of Minkowski spaces - a survey. Part I*, Expositiones Math., **19** (2001), 97–142.
- [6] H. MARTINI, SENLIN WU, *Radial projections of bisectors in Minkowski spaces*, Extracta Math., **23** (2008), 7–28.

- [7] H. MARTINI, SENLIN WU, *Halving circular arcs in normed planes*, Period. Math. Hungar., **57** (2008), 207–215.
- [8] A. C. THOMPSON, Minkowski Geometry, Encyclopedia of Mathematics and Its Applications, Vol. 63, Cambridge University Press, Cambridge, 1996.